

Prevention and treatment of acute kidney injury after cardiac surgery

Bengt Redfors

Department of Anaesthesiology and Intensive Care, Institute of Clinical Sciences, The Sahlgrenska Academy, University of Gothenburg, Sweden

Abstract

Acute kidney injury (AKI) occurs frequently after cardiac surgery and is independently associated with increased mortality. The main cause of AKI in these patients is renal ischemia. However, data on the renal oxygenation, defined as the renal oxygen supply/demand relationship are lacking in clinical AKI, and the effects of various pharmacological interventions on renal oxygenation are not known.

Patients and methods: The effects of mannitol (n=10) and dopamine (n=12) on renal blood flow (RBF), glomerular filtration rate (GFR) and renal oxygenation were analysed in post-cardiac surgery patients using the renal vein thermodilution technique. Furthermore, RBF, GFR and renal oxygenation were studied in patients with AKI (n=12) and compared to postoperative patients with no renal impairment (n=37). Finally, the effects of norepinephrine-induced changes in mean arterial pressure (MAP) on renal variables were analysed in AKI patients (n=12) with vasodilatory shock.

Results: Mannitol increased GFR and the renal oxygen demand (RVO₂), while it had no effect on RBF. Mannitol, thus, pharmacologically improved the renal function at the cost of an impaired renal oxygenation. In contrast, dopamine redistributed blood flow to the kidney and increased RBF, but had no effect on GFR or RVO₂. Consequently, dopamine improved renal oxygenation. AKI patients had a 40% lower RBF and a 60 % lower net-sodium reabsorption and GFR compared to control patients. However, contrary to previous hypothesis, this decrease in reabsorptive workload was not accompanied with a decrease in RVO₂. Thus, renal oxygenation was severely impaired in AKI. The high RVO₂ correlated directly to the sodium reabsorption, consuming 2.4 times more oxygen for a certain amount of reabsorbed sodium in AKI compared to control. Restoration of MAP from 60–75 mmHg with norepinephrine, improved renal oxygen delivery, GFR and renal oxygenation in AKI patients. Increasing MAP to 90 mmHg had no further beneficial effect.

Conclusions: While mannitol improves GFR at the cost of an impaired renal oxygenation, dopamine, in contrast, improves renal oxygenation, but has no effect on GFR. Furthermore, renal oxygenation is severely impaired in AKI, due to renal vasoconstriction and sodium reabsorption at a high oxygen cost. Finally, norepinephrine improves GFR and renal oxygenation when used for treatment of hypotension.

Key words: Kidney failure, acute; glomerular filtration rate; renal circulation; oxygen consumption; cardiac surgery; mannitol; dopamine; norepinephrine; autoregulation.

ISBN 978-91-628-8081-1 <http://hdl.handle.net/2077/21935>

Prevention and treatment of acute kidney injury after cardiac surgery

Akademisk avhandling

som för avläggande av medicine doktorexamen vid Sahlgrenska Akademien
vid Göteborgs Universitet kommer att offentligen försvaras i
Hjärtats aula, Sahlgrenska Universitetssjukhuset/Sahlgrenska,
onsdagen den 12 maj 2010, kl 09.00

av

Bengt Redfors
Leg. läkare

Fakultetsopponent:

Professor Sten Walther
Institutionen för medicin och hälsa
Universitetssjukhuset i Linköping, Linköping, Sverige

Avhandlingen baseras på följande delarbeten:

- I Redfors B, Sward K, Sellgren J, Ricksten SE
Effects of mannitol alone and mannitol plus furosemide on renal
oxygen consumption, blood flow and glomerular filtration after
cardiac surgery.
Intensive Care Med 2009, 35(1):115-122.
- II Redfors B, Bragadottir G, Sellgren J, Sward K, Ricksten SE
Dopamine increases renal oxygenation: a clinical study in post-
cardiac surgery patients.
Acta Anaesthesiol Scand, 54(2):183-190.
- III Redfors B, Bragadottir G, Sellgren J, Sward K, Ricksten SE
Acute renal failure is NOT an "acute renal success" – a clinical study on
renal oxygen supply/demand in postoperative acute kidney injury
Submitted, Critical Care Med
- IV Redfors B, Bragadottir G, Sellgren J, Sward K, Ricksten SE
Blood pressure restoration with norepinephrine improves renal
function and oxygenation in post-cardiac surgery patients with
vasodilatory shock and acute kidney injury
In manuscript

Göteborg 2010