



Det här verket har digitaliserats vid Göteborgs universitetsbibliotek och är fritt att använda. Alla tryckta texter är OCR-tolkade till maskinläsbar text. Det betyder att du kan söka och kopiera texten från dokumentet. Vissa äldre dokument med dåligt tryck kan vara svåra att OCR-tolka korrekt vilket medför att den OCR-tolkade texten kan innehålla fel och därför bör man visuellt jämföra med verkets bilder för att avgöra vad som är riktigt.

This work has been digitized at Gothenburg University Library and is free to use. All printed texts have been OCR-processed and converted to machine readable text. This means that you can search and copy text from the document. Some early printed books are hard to OCR-process correctly and the text may contain errors, so one should always visually compare it with the images to determine what is correct.



Ödsmål, Kville sn, Bohuslän

Hällristning  
Fiskare från  
bronsåldern

Rock carving  
Bronze age  
fishermen



MEDDELANDE från  
HAVSFISKELABORATORIET • LYSEKIL

nr  
254

Report of the Herring Otolith  
Workshop held at Lysekil  
18-19 September 1979

by

Rutger Rosenberg and R.J. Wood

October 1979

REPORT OF THE HERRING OTOLITH WORKSHOP HELD AT LYSEKIL 18-19 SEPTEMBER 1979

An otolith workshop was held at Lysekil (Sweden) in accordance with the following recommendation of the Herring Assessment Working Group for the area south of 62°N (ICES C.M. 1979/H:6 PELAGIC FISH COMMITTEE).

The Working Group recommends that otolith samples be collected from both spawning and juvenile herring and a workshop set up at an early date and convened by Dr. R. Rosenberg in Lysekil (Sweden) in order to determine the range of otolith types present in the indigenous stocks, to compare these with otoliths of the North Sea autumn spawners and to report to the 67th Statutory Meeting.

The following participated in the workshop

Dan Andreasson	Lysekil, Sweden	
Gunnar Aneer	Stockholm, Sweden	
Ad Corten	IJmuiden, Netherlands	
Oddvar Dahl	Bergen, Norway	
Didrik Danielssen	Arendal, Norway	
Bjarke Gloerfeldt-Tarp	Charlottenlund, Denmark	
Olle Hagström	Lysekil, Sweden	
Britt-Maj Karlsson	Lysekil, Sweden	
Niels Lundgaard	Charlottenlund, Denmark	
Marianne Martinsson	Lysekil, Sweden	
Rutger Rosenberg	Lysekil, Sweden	} JOINT CHAIRMEN
Jim Wood	Lowestoft, England	

A general discussion was held first of all on differences between the otoliths of spring and autumn spawning herring. Those considered were the appearance and size of the nucleus, the outline and shape of the whole otolith and the combined size of both the first growth and winter zones. It was decided, in view of the short length of time available, that the most profitable approach would be to attempt to separate samples of juvenile herring in Division IIIa into spring and autumn spawned fish on the basis of otolith differences in the size of the combined first growth and winter zones measured along the axis rostrum - post rostrum.

It was therefore necessary as a first step to examine the samples of otoliths which were available to the workshop from spawning herring of known origin. These consisted of spring spawners from:

The Kattegat: Randers Fjord (Denmark), Swedish coast and mid Kattegat.

The Skagerrak: Swedish coast and Norwegian fjords.

The Baltic: Central Swedish coast.

Autumn spawners from:

The Kattegat: Kobbergrund

The Baltic: South coast of Sweden

The North Sea: Orkney - Shetland, Longstone and Whitby

West of Scotland: Minch

The participants, working in small groups, were able to reach general agreement that the autumn spawners tended to have larger first zones than spring spawners and there was also a satisfactory agreement between measurements made on the same otoliths by the different groups. The results of the measurements made on the various samples of spawners are summarized in Table 1. It is clear that there is an increase in size across the table, northern Baltic spring spawners having the smallest and Orkney-Shetland autumn spawners the highest values. Tests of significance were carried out on mean sample values and the results are presented in Table 2. In most cases a significant difference was obtained between means and in only one of the comparisons between spring and autumn spawning fish could such a difference not be demonstrated.

An examination was then made of the otoliths which were available from juvenile herring caught in both the Skagerrak and Kattegat. The results are shown in Tables 3a and 3b. While the Kattegat samples seem to consist very predominantly of spring spawners, with a possible admixture of Baltic autumn spawners, those from the Skagerrak appear to include an autumn spawning component, particularly amongst the 1 ring fish, having a large first zone more characteristic of the North Sea herring.

On the basis of the results presented in this report it is recommended that measurements of otoliths should be continued in conjunction with counts of VS and  $K_2$ , especially in the Skagerrak, as an additional aid in separation of herring stocks in that area. It is suggested that this type of examination could prove of value in many other areas.

This paper was presented at the 67th Statutory Meeting in Warszawa, Poland, in October 1979. Reference number C.M.1979/H:66.

Table 1 Percentage distribution of otolith first zone measurements in samples of spawning herring from various localities.

* epu's	Spring spawners			Autumn spawners			I Orkney/ Shetland			
	A Northern Baltic - Askö area	B Norwegian coast	C Danish coast - Randers fjord	D Swedish coast	E South- western Baltic	F Kattegat - Kobbergrund		G West of Scotland North Sea	H Central North Sea	
30-34	2.0	1.7								
35-39	20.4	8.0								
40-44	<u>42.9</u>	4.5								
45-49	28.6	13.6	3.8						1.0	
50-54	4.1	14.2	5.0	2.4	4.1	0.5			2.0	
55-59	2.0	<u>24.4</u>	28.7	9.5	4.2	3.7			4.1	
60-64		19.9	<u>30.0</u>	30.9	16.7	6.4			2.0	
65-69		7.4	28.7	<u>35.7</u>	<u>54.2</u>	18.7			8.2	2.4
70-74		6.3	3.8	16.7	12.5	<u>32.7</u>			14.3	16.7
75-79				4.8	8.3	24.6			<u>27.6</u>	28.5
80-84						9.1			26.5	<u>33.3</u>
85-89						3.2			11.8	16.7
90-94						1.1			2.0	2.4
									1.0	
Mean	45.31	56.60	63.65	67.44	67.83	70.24	70.00	73.97	76.90	
(No mea- sured)	(49)	(176)	(80)	(42)	(24)	(187)	(106)	(98)	(42)	
Variance	22.87	81.04	31.84	30.64	34.40	47.66	25.49	85.78	33.28	
Mean VS	55.29	57.12	55.75	56.54	55.56	56.18	56.10	56.39	56.45	
Mean K <sub>2</sub>	13.40	-	13.67	13.88	-	14.15	14.27	14.26	14.11	

\* 25 epu's = 1 mm

Table 2.

Test of significance between means of otolith measurements given in Table 1. Values of "t" in Student's t-test are given along with the degree of significance. \*\*\* 0.001, \*\* 0.01, \* 0.05; N.S. indicates not significant.

	A	B	C	D	E	F	G	H	I
A		***	***	***	***	***	***	***	***
B	11.7		***	***	***	***	***	***	***
C	19.7	7.6		***	**	***	***	***	***
D	20.2	9.9	3.6		N.S.	*	*	***	***
E	16.3	8.2	3.1	0.3		N.S.	N.S.	***	***
F	29.4	16.0	7.9	2.6	1.7		N.S.	***	***
G	29.3	16.1	8.2	2.8	1.8	0.3		***	***
H	24.7	15.0	9.1	5.1	4.0	3.8	3.5		**
I	28.2	18.1	12.1	7.7	6.1	6.8	6.5	2.7	

Table 3(a) Juvenile herring from the Skagerrak

	1 ring N of Hirtshals 31-8-78	1 ring N of Hirtshals 11-2-76	1 ring N of Hirtshals 12-8-78	1 ring N of Hanst- holm 16-7-79	1 ring off Fredrikstad 1-8-77	1 ring 15 miles N of Skagen 19-6-78	1 ring North of Hanstholm 7-2-78
30-34							
35-39							
40-44							
45-49							
50-54	1	3		5			1
55-59	5	11	10	13	5	5	2
60-64	31	20	29	15	14	10	4
65-69	25	10	8	2	40	4	3
70-74	17	2	2	19	22	1	3
75-79	11	2	1	10	5	7	4
80-84	3	5	4	4		8	3
85-89	1	2				2	4
90-94	1						3
95-99						1	4
Mean	70.21	63.89	81.11	72.73	68.33	82.63	86.48
Variance	47.32	22.10	26.54	27.11	49.80	40.44	25.58
No. of fish	95	46	10	58	91	19	32
			50	35		19	20
Mean VS	56.28	56.34	56.00	56.57	-	55.78	-
Mean K <sub>2</sub>	14.01	-	-	-	-	14.00	-
			56.06	56.20	-	55.95	-
			-	-	-	13.68	-

continued

Table 3(a) continued

	1 ring NW of Hirtshals 12-8-78	1 ring N of Hirtshals 11-2-76	2 ring
30-34			
35-39			
40-44		1	
45-49	1		1
50-54	1		1
55-59	4	1	4
60-64	14	4	16
65-69	7	8	7
70-74	2	5	12
75-79	1	5	6
80-84	4	5	
85-89	2	4	
90-94			
95-99			
Mean	67.43	73.62	66.51
Variance	80.53	61.89	63.41
No. of fish	36	32	48
Mean $V_s$		56.18	56.34
Mean $K_2$		14.03	14.09



Table 3(b) Juvenile herring from the Kattegat.

	1 ring Schults ground 3-4-79	2 ring	1 ring North Kattegat 16-11-78	1 ring NE of Anholt 9-5-79	2 ring	1 ring Southern Kattegat 10-10-78	1 ring 10 miles west of Falkenberg 28-11-78	2 ring
30-34								
35-39								
40-44								
45-49				3				3
50-54	1	12	1	4	5	16	1	9
55-59	10	45	16	2	15	55	55	36
60-64	20	23	14	8	20	47	78	58
65-69	21	3	4	2	3	16	53	24
70-74	11	3	1	1		4	30	2
75-79	3	1					22	
80-84							7	
85-89							8	
90-94								
95-99								
Mean	67.19	61.38	62.64	59.75	61.59	62.26	67.98	63.36
Variance	29.59	20.51	17.34	51.19	15.17	19.16	46.91	23.83
No. of fish	66	87	36	20	43	128	254	132
Mean VS	55.92	55.88	56.22	55.60	55.72	55.97	56.18	55.88
Mean K <sub>2</sub>	13.83	13.81	-	-	-	13.92	13.94	13.97

