PERITONEAL FIBRINOLYSIS DURING

PNEUMOPERITONEUM AND LAPAROSCOPIC SURGERY

AKADEMISK AVHANDLING

Som för avläggande av medicine doktorsexamen vid Göteborgs Universitet kommer att offentligt försvaras I Bakfickan, Servicehuset, Sahlgrenska Universitetssjukhuset-Östra, Göteborg torsdagen 6 december 2007 13.00

av

MARIA BERGSTRÖM leg läkare

Avhandlingen baseras på följande delarbeten:

- I. Peritoneal response to pneumoperitoneum and laparoscopic surgery. Bergström M, Ivarsson ML, Holmdahl L, *Br J Surg 2002;89(11):1465-9*
- II. CO₂ promotes plasminogen activator inhibitor type 1 expression in human mesothelial cells
 Bergström M, Falk P, Holmdahl L, Surg Endosc 2003;17(11):1818-22
- III. Effect of acidosis on expression of mesothelial cell plasminogen activator inhibitor type-1 Bergstrom M, Falk P, Holmdahl L, Surg Endosc 2006;20(9):1448-52
- IV. Peritoneal and systemic pH during Pneumoperitoneum with CO₂ and helium in a pig model
 Bergstrom M, Falk P, Park PO, Holmdahl L,
 Surg Endosc 2007, e-pub ahead of print
- V. Clinical evaluation of peritoneal acidification and fibrinolytic response during laparoscopy, a randomized parallel group study comparing Helium and Carbon dioxide Bergstrom M, Falk P, Park PO, Bengtsson JP, Haglind E, Holmdahl L, *in manuscript*

PERITONEAL FIBRINOLYSIS DURING PNEUMOPERITONEUM AND LAPAROSCOPIC SURGERY

Maria Bergström, dept of Surgery, Sahlgrenska University Hospital-Östra, University of Göteborg, S-416 85 Göteborg, Sweden

BACKGROUND

Laparoscopic surgery is believed to induce less postoperative adhesion formation compared with open procedures, but information regarding biological impact of a laparoscopic approach is limited.

MATERIAL & METHODS

Peritoneal response to laparoscopic surgery was assessed in human peritoneal tissue in two clinical trials (paper I & V). Human mesothelial cell response to CO_2 was assessed in cell culturing media (paper II & III). Measurements of peritoneal pH was evaluated in an animal model (paper IV), and then performed in patients (paper V). Assays of the key fibrinolytic enzymes t-PA and PAI-1 were done at protein (paper I-III & IV) and mRNA levels (paper II & III).

RESULTS

The initial clinical study showed a similar decrease in peritoneal t-PA activity during both open and laparoscopic cholecystectomy. However, there was a higher initial peritoneal PAI-1 concentration in the laparoscopic group, which might be attributable to prior exposure to CO_2 . The *in vitro* studies showed an up regulation of PAI-1 mRNA production in cells exposed to CO_2 or acidic conditions, and that acidification could be caused by CO_2 . *In vivo* studies of peritoneal pH showed an immediate decrease to 6.5 during insufflation, reproducing the CO_2 effect during surgery. He did not affect peritoneal pH. When initial peritoneal exposure to CO_2 was controlled for, peritoneal levels of t-PA decreased and PAI-1 increased during laparoscopic surgery regardless of gas used. These findings are consistent with observations done in open surgery and indicate that the effect is related to the surgical trauma. However, the t-PA activity was better preserved using CO_2 suggesting that use of CO_2 might have less adverse effect on peritoneal fibrinolysis than He.

CONCLUSION

Peritoneal fibrinolytic response during laparoscopic surgery is similar to open surgery, but CO_2 elicits specific biological effects. Exposure of peritoneum and mesothelium to CO_2 leads to a local and systemic acidosis and seems to have direct effects on key fibrinolytic enzymes. The systemic acidosis is manageable through controlled ventilation. The clinical implication of the effect on peritoneal fibrinolysis is unclear, but the observations done in humans are consistent with a reduced propensity for adhesion formation.

Key words: Laparoscopy, Peritoneum, Fibrinolysis, Mesothelial cells, Carbon dioxide, Helium ISBN: 978-91-628-7368-4